

Universality of monopole mode and time evolution of a d -dimensional trapped interacting Bose gas

Tarun Kanti Ghosh

The Institute of Mathematical Sciences, C. I. T. Campus, Chennai-600 113, India.

(February 1, 2008)

We study a generalised Gross-Pitaevskii equation describing a d -dimensional harmonic trapped (with trap frequency ω_0) weakly interacting Bose gas with a non-linearity of order $(2k + 1)$ and scaling exponent (n) of the interaction potential. Using the time-dependent variational analysis, we explicitly show that for a particular combination of n , k and d , the generalised GP equation has the universal monopole oscillation frequency $2\omega_0$. We also find that the time-evolution of the width can be described universally by the same Hill's equation if the system satisfy that particular combination. We also obtain the condition for the exact self-similar solutions of the Gross-Pitaevskii equation. As an application, we discuss low dimensional trapped Bose condensate state and Calogero model.

PACS numbers:05.45.-a, 03.75.Fi.

I. INTRODUCTION

There has been renewed interest in the Bose-Einstein condensation (BEC) after its experimental achievement by several groups [1]. The Gross-Pitaevskii (GP) equation explains most of the properties of these trapped dilute interacting Bose particles [2]. After the discovery of BEC in a trapped alkali atom, the influences of the dimension of a Bose system is being the subject of extensive study [3–7]. Recently, low dimensional BEC has been observed in MIT [8]. It is well known that the nonlinear GP equation for a 2D trapped Bose system has a monopole mode with a universal frequency $2\omega_0$ [3,9,10]. In 2D trapped interacting Bose system, the Fermi pseudopotential interaction does not affect the monopole mode frequency. This peculiar behavior of monopole mode frequency depends on the scaling property of the interaction potential and the order of non-linearity. Without trap potential, the GP equation is scale invariant. Introduction of harmonic potential breaks the scale invariance. Pitaevskii and Rosch [3] has pointed out that due to special property of the harmonic potential, SO(2,1) symmetry still exists. They have shown the universal nature of the monopole mode frequency only for 2D GP equation by constructing the generators of the SO(2,1) symmetry [3]. Here, using the time-dependent variational approach, we obtain a condition for the universality of monopole mode frequency and the dynamics of width of a class of GP equation describing the trapped interacting Bose system, at varying spatial dimensionality, order of the nonlinearity and the scaling exponent of the interaction potential. Interestingly, the dynamics of the width of a class of GP equation can be described by the same Hill's equation if the Bose system satisfy that particular combination or shows the universal monopole mode frequency.

This paper is organised as follows. In Sec. II, using the time-dependent variational analysis, we show that for a particular combination of n , k and d the generalised GP equation describing the d -dimensional harmonic trapped Bose gas exhibits the universal monopole oscillation frequency $2\omega_0$. We also derive the time-evolution of the width of the d -dimensional trapped Bose system which satisfies that particular combination. The time evolution of the width can be described universally by the same Hill's equation which is analytically solvable. We also discuss the condition for exact self-similar solutions of the GP equation. In Sec.III, we consider quasi-2D trapped Bose gas interacting through the Fermi pseudopotential which can be described by an *ordinary* GP equation. In Sec. IV, we also consider the 1D *modified* GP equation which describes the BEC state. We obtain the universal monopole mode frequency and time evolution of the condensate width of this system in a time-independent trap as well as a time-dependent trap. In Sec. V, we discuss the Calogero type potential. In Sec. VI, we present the summary and conclusions of our work.

II. TIME-DEPENDENT VARIATIONAL ANALYSIS

To show the universality of the monopole mode frequency and the dynamics of the width of a class of GP equation (also known as nonlinear Schrodinger equation (NLSE)), we use the time dependent variational method. This variational method has been extensively and successfully used to explain many properties in BEC and other nonlinear systems. Using this method, Garcia *et al.* [11] derived a general dispersion relation for the low energy excitations of a 3D deformed trapped Bose gas which exactly coincides with the results obtained by Stringari [12] for large number of particles.

Consider a system of Bose particles in d -dimensional harmonic trapped potential $V_t(\vec{r}) = \frac{1}{2}M\omega_0^2 r^2$ at zero temperature and interacting with a translationally invariant interaction potential $V_I = g_d V(\vec{r} - \vec{r}')$, where $r^2 = \sum_{i=1}^d x_i^2$ and g_d is the coupling constant in d dimension. M is the atomic mass and $\vec{r}$ is a d -dimensional vector. Under the scale transformation, $\vec{r} \rightarrow \lambda \vec{r}$, we assume $V_I(\lambda \vec{r}) = \lambda^n V_I(\vec{r})$ where n is the scaling exponent of the interaction potential.

We are considering a generalized d -dimensional GP equation [13] describing a d -dimensional harmonic trapped weakly interacting Bose gas with a non-linearity of order $(2k+1)$ and scaling exponent n of an arbitrary interaction potential $V_I = g_d V(\vec{r} - \vec{r}')$

$$i\hbar \frac{\partial \psi(\vec{r})}{\partial t} = [-\frac{\hbar^2}{2M} \nabla^2 + V_t(r) + g_d \int d^m r' \psi^*(\vec{r}') V(\vec{r} - \vec{r}') \psi(\vec{r}') |\psi(\vec{r})|^{2(k-1)}] \psi(\vec{r}), \quad (1)$$

where ∇^2 is the d -dimensional Laplacian operator and $*$ denotes complex conjugation. The generalised d -dimensional energy functional is

$$E[\psi] = \int d^d r [\frac{\hbar^2}{2M} |\nabla \psi|^2 + V_t |\psi|^2 + \frac{g_d}{k+1} \int d^m r' \psi^*(\vec{r}) \psi^*(\vec{r}') V(\vec{r} - \vec{r}') \psi(\vec{r}') |\psi(\vec{r})|^{2(k-1)} \psi(\vec{r})]. \quad (2)$$

If $m = d =$ arbitrary dimension and $V(\vec{r} - \vec{r}')$ is a d -dimensional delta function potential, the above equation gives the non-linear term of order $(2k+2)$. If $k = 1$ and $m = d \geq 1$, Eq. (1) becomes *ordinary* GP equation. and if $k = 2$ and $m = d = 1$, Eq. (1) gives the 1D *modified* GP equation in which the $|\psi|^6$ interaction term is present [14,15].

One can write down the generalised Lagrangian density corresponding to this system which is

$$L = \frac{i\hbar}{2} (\psi \frac{\partial \psi^*}{\partial t} - \psi^* \frac{\partial \psi}{\partial t}) + [\frac{\hbar^2}{2M} |\nabla \psi|^2 + V_t(r) |\psi(\vec{r})|^2 + \frac{g_d}{(k+1)} \int d^m r' \psi^*(\vec{r}) \psi^*(\vec{r}') V(\vec{r} - \vec{r}') \psi(\vec{r}') |\psi(\vec{r})|^{2(k-1)} \psi(\vec{r})]. \quad (3)$$

Without interaction potential, Eq. (1) reduces to the linear Schrodinger equation which has the Gaussian wave function for the ground state. In order to get the evolution of the square of the radius, we assume the following Gaussian wave function,

$$\psi(r) = C(t) e^{-\frac{r^2}{2a_0^2} [\frac{1}{\alpha(t)} + i\beta(t)]}, \quad (4)$$

where $C(t)$ is the normalization constant which can be determined by the normalization condition $\int |\psi|^2 d^d r = N$ and $a_0 = \sqrt{\hbar/M\omega_0}$ is the oscillator length. N is the total number of particles in the system. $\alpha(t)$ and $\beta(t)$ are the time-dependent dimensionless variational parameters. $\alpha(t)$ is the square of the radius of the system. In order to describe the dynamics of the variational parameter $\alpha(t)$, the phase factor $i\beta(t)r^2$ is required [11]. Note that the same kind of phase factor is also present in the Eq. (4) of the Ref. [10] to get the dynamics of the width of a system. We get the effective Lagrangian $L_{eff}(\alpha, \beta)$ by substituting Eq. (4) into Eq. (3) and integrating the Lagrangian density over the space coordinates,

$$L_{eff} = \frac{N\hbar\omega_0}{4} [-\alpha\dot{\beta}d + \frac{d}{\alpha} + \alpha d(1 + \beta^2) + G_d \alpha^{(n+m-kd)/2}], \quad (5)$$

where

$$G_d = G_0 \int d^d R d^m R' e^{-(R^2 + R'^2)} V(\vec{R} - \vec{R}') e^{-(R'^2 + R^2(2k-1))}, \quad (6)$$

where $G_0 = \frac{4g_d N^k 2^{(d+m+n)/2} a_0^{(n+m-kd)}}{(k+1)\hbar\omega_0\pi^{(k+1)d/2}}$ and $r^2 = 2\alpha R^2 a_0^2$.

Using the Lagrange equation of motion, we get the equation of motion for the variational parameter $\alpha(t)$. The time evolution of the variational parameter $\alpha(t)$ is

$$\ddot{\alpha} = 2\alpha\beta^2 + \frac{2}{\alpha} - 2\alpha - \frac{(n+m-kd)G_d}{d}\alpha^{(n+m-kd)/2}. \quad (7)$$

The "." represents the derivative with respect to the dimensionless variable $\tau = \omega_0 t$. The total energy of the system is

$$\tilde{E} = 4E/N\hbar\omega_0 = \alpha\beta^2 d + \frac{d}{\alpha} + \alpha d + G_d \alpha^{(n+m-kd)/2}. \quad (8)$$

At equilibrium, $\beta = 0$ and $\frac{\partial E}{\partial \alpha}|_{\alpha=\alpha_0} = 0$. The equilibrium point α_0 satisfies the following relation in d -dimension

$$\alpha_0^2 + \frac{(n+m-kd)G_d}{2d}\alpha_0^{(n+m-kd)+1} = 1. \quad (9)$$

We choose n , m , k and d such that they satisfy the following relation

$$n + m - kd + 2 = 0. \quad (10)$$

If n , m , k , and d satisfy the above relation, Eq. (7) can be written as

$$\ddot{\alpha} + 4\alpha = 2\tilde{E}(\alpha). \quad (11)$$

Define $I(t) = \int d^d r |\psi(\vec{r}, t)|^2 r^2$ which is the expectation value of the square of the radius. The above equation can be re-written as

$$\ddot{I} + 4\omega_0^2 I = \frac{4E(I)}{M}. \quad (12)$$

From now onwards "." indicates derivative with respect to the time t . For monopole excitation, the average value of the collective coordinate r^2 oscillates around its equilibrium value. We expand $I(t)$ around its equilibrium point I_0 as $I(t) = I_0 + \delta I(t)$ and we get,

$$\delta \ddot{I} + 4\omega_0^2 \delta I = \frac{4E_0}{M}, \quad (13)$$

where $E_0 = d(\frac{1}{\alpha_0} + \alpha_0 + \frac{G_d}{d})$ is the equilibrium energy of the system. This equation is identical with the classical equation (1) of Ref. [3] which was derived using the virial theorem. The solution of this equation is

$$I(t) = A \cos(2\omega_0 t + \theta) + \frac{E_0}{M\omega_0^2}. \quad (14)$$

A and θ are constants which can be determined from the initial conditions on ψ . Thus the square of the system radius oscillates with a frequency $2\omega_0$ in any dimension if $n + m - kd + 2 = 0$. This monopole frequency $2\omega_0$ is universal because it is independent of the interatomic interaction, dimensionality of the trap, order of non-linearity and the total number of atoms N . For *ordinary* GP equation in $d(=m)$ dimension, $k = 1$. Hence the monopole

mode frequency of an *ordinary* d -dimensional GP equation is the universal frequency $2\omega_0$ if $n = -2$. Eqs. (7) and (10) explicitly shows how the scaling exponent of the interaction potential and the order of non-linearity gives the universal monopole mode frequency.

If n, m, d and k satisfy the Eq. (10), the equation (7) can be re-written for time-independent trap as well as for time-dependent trap as

$$\ddot{I}(t) - \frac{\dot{I}^2(t)}{2I(t)} + 2\omega^2(t)I(t) = \frac{Q_d}{I(t)}, \quad (15)$$

where $Q_d = (\frac{N\hbar d}{2M})^2(1 + G_d)$ is the positive invariants under time evolution. This Q_d does not depend on the frequency of the trap potential. Let $X(t) = \sqrt{I(t)}$ is the condensate width. The above equation can be written as

$$\ddot{X}(t) + \omega_0^2(t)X(t) = \frac{Q_d}{X^3(t)}. \quad (16)$$

This is a non-linear singular Hill's equation in d dimension. This is an interesting result because the time evolution of the width can be described by the same Hill's equation in any dimension if $n + m - kd + 2 = 0$. The dynamics of the width is universal because the time evolution of the width can be described universally by the same non-linear singular Hill's equation in any dimension. Moreover, this Hill's equation can be solved analytically and exactly. The Eq.(16) can be viewed as the classical motion of a fictitious particle in an effective d -dimensional potential

$$V_{eff} = \frac{\omega_0^2(t)X^2}{2} + \frac{Q_d}{2X^2}. \quad (17)$$

The general solution of Eq.(16) is $X(t) = \sqrt{u^2(t) + \frac{Q_d}{W^2}v^2(t)}$ where $u(t)$ and $v(t)$ are two linearly independent solutions of the equation $\ddot{p} + \omega^2(t)p = 0$ which satisfy the initial conditions $u(t_0) = X(t_0)$, $\dot{u}(t_0) = X'(t_0)$, $v(t_0) = 0$ and $v'(t_0) \neq 0$. W is the Wronskian. This time-dependent Hill's equation can be solved explicitly only for particular choices of $\omega_0(t)$. For time-dependent periodic trap potential, the equation is known as Mathieu equation which is well studied. In most of the BEC experiment, the trap potential is time-independent. For time-independent trap potential, a solution of this equation is

$$X(t) = \sqrt{\cos^2(2\omega_0 t) + \frac{Q_d}{4\omega_0^2} \sin^2(2\omega_0 t)}. \quad (18)$$

From Eq.(10), we get $kd = 2$ for 1D *modified* GP equation, and 2D *ordinary* GP equation with $n = -2$. This is the condition for self-similar solution of a non-linear GP equation discussed in by Rybin [14,15]. There is no self-similar solution in 1D *ordinary* GP equation since it does not obey Eq.(10).

We now discuss few systems which satisfy the Eq. (10) and shows the universal monopole mode frequency and the dynamics of the width of the system that can be described by the Eq. (18).

III. 2D TRAPPED BOSE GAS

It has been shown by Baganato *et al.* [16] that for an ideal 2D Bose gas under harmonic trap a macroscopic occupation of the ground state can exist at temperature $T < T_c = \sqrt{N/\zeta(2)}\hbar\omega/k_B$. With present technology one can freeze the motion of the trapped particles in one direction to create a quasi-2D Bose gas. In the frozen direction the particles execute the zero point motion. To achieve this quasi-2D system, the frequency in the frozen direction should be much larger than the frequency in the X-Y plane and the mean interactions between the particles. The wave function in the z-direction is

$$\psi(z) = Ce^{-\frac{z^2}{2a_z^2}}, \quad (19)$$

where $a_z = \sqrt{\hbar/m\omega_z}$ is the oscillator length in the z-direction and C is the normalization constant. When we integrate out the z-component in the 3D GP energy functional, then we get the effective interaction potential $V_I = g_2\delta^2(\vec{r} - \vec{r}')$ [7] where $g_2 = 2\sqrt{2\pi}\hbar\omega_z aa_z$ is the effective coupling strength in 2D. a is the s -wave scattering length in 3D. This quasi-2D BEC state can be described by the *ordinary* GP equation which is valid only when $a < a_z$ [5]. The same effective coupling constant is obtained in [4]. Under the scale transformation, $\vec{r} \rightarrow \lambda\vec{r}$, $V_I(\lambda\vec{r}) = \frac{1}{\lambda^2}V_I(\vec{r})$. So this interaction potential has the scaling exponent $n = -2$. We will show that for an isotropic harmonic trapped system, the monopole mode frequency is $2\omega_0$. In this system, $m = d = 2$ and $k = 1$ which satisfy the Eq. (10).

The time-evolution of this system is

$$\ddot{X} + \omega_0^2 X = \frac{Q_2}{X^3}. \quad (20)$$

The same equation is obtained by using different method at zero temperature [10,17]. Here, we have identified the value of Q_2 at $T = 0$ which is $Q_2 = \frac{N^2\hbar^2}{M^2}(1 + \sqrt{\frac{2}{\pi}}N\frac{a}{a_z})$. The dynamics of the width can be described by the Eq. (18). At equilibrium, $X_0^4 = \frac{Q_2}{\omega_0^2}$. When we linearize Eq. (20) around the equilibrium point X_0 , we get,

$$\delta\ddot{X} + 4\omega_0^2\delta X = 0. \quad (21)$$

Once again we obtain the universal oscillation frequency of the condensate width to be $\omega = 2\omega_0$, corresponding to the frequency of a single particle excitation in the condensate state.

IV. 1D TRAPPED BOSE GAS

It has been shown by Kolomeisky *et al.* [15] that the GP equation describing BEC in 1D contains $|\psi|^6$ interaction term instead of familiar $|\psi|^4$ term. The energy functional of 1D dilute interacting Bose gas at zero temperature is

$$E = \int dx \left[\frac{\hbar^2}{2M} \left| \frac{d\psi}{dx} \right|^2 + V(x)|\psi|^2 + \frac{\hbar^2\pi^2}{6M} |\psi|^6 \right]. \quad (22)$$

In this system, $d = m = 1$, $k = 2$ which satisfy the Eq. (10) and $g_1 = \frac{\hbar^2\pi^2}{2M}$. The time-evolution of the system is

$$\ddot{X} + \omega_0^2 X = \frac{Q_1}{X^3}. \quad (23)$$

where $Q_1 = \frac{N^2\hbar^2}{4M^2}[1 + \frac{4\pi N^2}{3\sqrt{3}}]$. The same equation is obtained in [14,15] by using different method. But here we have identified the value of constant Q_1 . This 1D *modified* GP equation also exhibits an universal monopole mode frequency $2\omega_0$ and satisfy the self-similar condition. The dynamics can be described by the Hill's equation which is exactly solvable.

For 1D ordinary GP equation, $m = d = k = 1$, which does not satisfy the relation (10). So the monopole mode frequency is affected by the interatomic strength and number of particles and the time-evolution of the system can not be described by non-linear singular Hill's equation.

V. CALOGERO MODEL

Only an inverse square pair potential, $1/r^2$, has the scaling, $n = -2$, in any dimension and it satisfies the Eq. (10) for $k = 1$ in arbitrary d . A system of trapped particles interacting through pair potential $1/r^2$ is well known as the Calogero-Sutherland model [18]. In Ref. [19], they have shown that the energy spectrum of a system of harmonic trapped particles interacting with $1/r^2$ potential is divided into sets of equidistant levels with separation $2\omega_0$ in any dimension. In Ref. [20], they have considered a one-dimensional system of particles interacting by an inverse square pair potential and shown that the monopole mode frequency is the universal frequency $2\omega_0$. For $m = k = 0$ and $n = -2$, we get Calogero model [18] in arbitrary d which gives the same universal monopole mode frequency $2\omega_0$. In this system, the dynamics of the width can be described universally by the Eq. (18).

Apart from these two potentials, there are some other model potentials which have the scaling exponent $n = -2$. For example, the derivative delta function potential in 1D, the double delta function potential in 1D, and the $1/x$ delta function potential in 1D. For all these model potentials, one can get the universal monopole mode frequency and the dynamics of the width.

VI. SUMMARY AND CONCLUSIONS

In this work, using the time-dependent variational analysis, we have obtained the condition, $n + m - kd + 2 = 0$, for which a class of NLSE exhibits the universal monopole oscillation frequency $2\omega_o$. This monopole mode frequency is universal because it does not depend on the strength of the interaction potential, dimensionality of the trap, order of non-linearity and total number of atoms N . We have also shown the universality of time evolution of the width in a time-independent trap as well as a time-dependent trap for a class of GP equation. Interestingly, this dynamics of the width can be described by the same Hill's equation (16) if the Bose system satisfy the condition, $n + m - kd + 2 = 0$. This Hill's equation is analytically and exactly solvable. Using the time dependent variational analysis, we have also obtained the condition for self-similar solutions, $kd = 2$, of a class of non-linear GP equation.

As an example, we have shown that the 2D trapped Bose gas describing by the *ordinary* GP equation, 1D BEC describing by the *modified* GP equation and a system describing by the Calogero model shows the universal nature of the monopole mode frequency and the dynamics of the width of the system. We have identified the values of the positive invariants quantity, Q_2 and Q_1 for 2D and 1D BEC.

After submitting this paper on the net (cond-mat/0012188), there is another paper [21] appeared where the universality of the monopole mode and the dynamics of the width of a class of NLSE has been shown exactly and analytically which is obtained from different method. Note that the phase factor in the variational ansatz of the wave function (eq.(4)) is required to get the dynamics of the width [11]. Similarly, the phase factor is also present in equation of the time dependent transformations in [10,21]. In Ref. [21], using those time dependent transformations, the universality of the monopole mode and the dynamics of the width is obtained for a class of NLSE. Our results obtained from the variational approach are identical to the exact results of [21]. This is because of the similarity between the variational ansatz for the wave function and the transformation equation (eq.(2)) in Ref. [21].

Our result is also valid for $SO(2,1)$ invariant multicomponent NLSE describing BEC state. Recently, 2D and 1D BEC has been observed in MIT [8]. Experimentally, it is also possible to check the validity of the two-body potential in 2D BEC state [3] and the order of the nonlinearity in 1D BEC state by measuring the universal monopole mode frequency and the dynamics of the width of the condensate state.

I would like to thank Pijush K. Ghosh, M. V. N. Murthy and K. Santosh for helpful discussions and many valuable suggestions.

- [1] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell, *Science*, 269, (1995) 198; C. C. Bradley, C. A. Sackett, J. J. Tollet, and R. G. Hulet, *Phys. Rev. Lett.* 75, (1995), 1687; K. B. Davis, M. O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle, *Phys. Rev. Lett.* 75, (1995), 3969.
- [2] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, *Reviews of Modern Physics*, **71**, 463 (1999); A. S. Parkins and D. F. Walls, *Phys. Rep.* 303, (1998), 1; A. Griffin, Preprint, cond-mat/ 9911419.
- [3] L. P. Pitaevskii and A. Rosch, *Phys. Rev. A*, 55, (1997), R853.
- [4] Tin-Lun Ho and Michael Ma, *J. Low Temp. Phys.* 115, (1999), 61.
- [5] D. S. Petrov, M. Holzmann and G. V. Shlyapnikov, *Phys. Rev. Lett.* 84, (2000), 2551.
- [6] Tarun K. Ghosh, *Phys. Rev. A*, 63, (2001), 013603.
- [7] Tarun K. Ghosh and Subhasis Sinha, cond-mat/0102258.
- [8] A. Gorlitz *et al.* cond-mat/0104549.
- [9] L. P. Pitaevskii, *Phys. Lett. A*, 221, (1996), 14.
- [10] Yu. Kagan, E. L. Sukrov, and G. V. Shlyapnikov, *Phys. Rev. A*, 54, (1996), R1753.
- [11] Victor M. Perez-Garcia, H. Michinel, J. I. Cirac, M. Lewenstein, and P. Zoller, *Phys. Rev. Lett.* 77, (1996), 5320.
- [12] S. Stringari, *Phys. Rev. Lett.* 77, (1996), 2360.
- [13] E. P. Gross, *Nuovo Cimento A*, 20, (1961), 454; L. P. Pitaevskii, *Pis'ma Zh. Eksp. Teor. Fiz.* 77, (1961), 988 [*Sov. Phys. JETP*. 13, (1961), 451].
- [14] A. Rybin, G. Varzugin, M. Lindberg, J. Timonen, R. K. Bullough, cond-mat/0001059.
- [15] E. B. Kolomeisky, T. J. Newman, Joseph, Joseph P. Straley, Xiaoya Qi, *Phys. Rev. Lett.* 85, (2000), 1146.
- [16] V. Baganato and D. Kleppner, *Phys. Rev. A*, 44, (1991), 7439.
- [17] Juan J. Garcia-Ripoll and Victor M. Perez-Garcia, *Phys. Rev. Lett.* 83, (1999), 1715.
- [18] F. Calogero, *J. math. Phys. (N. Y.)* 10, (1969), 2191; 10, (1969), 2197; B. Sutherland, *ibid.* 12, (1971), 246; 12, (1971), 251.
- [19] S. C. Benjamin, L. Quiroga, and N. F. Johnson, *Phys. Rev. A*, 54, (1996), 4309.
- [20] B. Sutherland, *Phys. Rev. Lett.* 80, (1998), 3678.
- [21] Pijush K. Ghosh, cond-mat/0102488